

# **Lanesend Primary School**

## **Diabetes Policy 2023** School Health and Safety Policy

| Signed:       | Date: |
|---------------|-------|
| (Headteacher) |       |

Signed: ..... Date: (Chair of Trustees)

Review Date: April 2024 Reviewed By: Headteacher and Child-Centred Group

#### Lanesend Primary Diabetes Policy

#### What is Diabetes?

- Diabetes is a condition where the level of glucose in the blood rises. This is either due to the lack of insulin (Type 1 diabetes) or because there is insufficient insulin for the child's needs or the insulin is not working properly (Type 2 diabetes).
- The majority have Type 1 diabetes. They normally need to have daily insulin injections, to monitor their blood glucose level and to eat regularly according to their personal dietary plan. People with Type 2 diabetes are usually treated by diet and exercise alone.
- Each person may experience different symptoms and this should be discussed when drawing up an individual health and risk assessment. Greater than usual need to go to the toilet or to drink, tiredness and weight loss may indicate poor diabetic control, and staff will naturally wish to draw any such signs to the families' attention.
- Staff with diabetes should make their condition known and their treatment plan available.
- If a member of staff or a child has diabetes in their class, children and staff will use the emergency lanyards to get another adult should they become unwell. A diabetes information session will be shared with the class at the beginning of the year.

#### Medicine and Control for children

The diabetes of the majority of children is controlled by injections of insulin each day. Most younger children will be on a twice a day insulin regime of a longer acting insulin and it is unlikely that these will need to be given during school hours, although for those who do, it may be necessary for an adult to administer the injection. Older children may be on multiple injections and others may be controlled on an insulin pump. Most children can manage their own injections, but if doses are required at school supervision may be required, and also a suitable, private place to carry it out.

Increasingly, older children are taught to count their carbohydrate intake and adjust their insulin accordingly. This means that they have a daily dose of long-acting insulin at home, usually at bedtime; and then insulin with breakfast, lunch and the evening meal, and before substantial snacks. The child is taught how much insulin to give with each meal, depending on the amount of carbohydrate eaten. They may or may not need to test blood sugar prior to the meal and to decide how much insulin to give. Diabetic specialists would only implement this type of regime when they were confident that the child was competent. The child is then responsible for the injections and the regime would be set out in the individual health and risk assessment.

Children with diabetes need to ensure that their blood glucose levels remain stable and may check their levels by taking a small sample of blood and using a small monitor at regular intervals. They may need to do this during the school lunch break, before PE or more regularly if their insulin needs adjusting. Most older children will be able to do this themselves and will simply need a suitable place to do so. However, younger children may need adult supervision to carry out the test and/or interpret test results.

When staff agree to administer blood glucose tests or insulin injections, they should be trained by an appropriate health professional. Administering injections is a matter for personal preference and no member of staff will be expected to carry out this task without full training and their consent.

Children with diabetes need to be allowed to eat regularly during the day. This may include eating snacks during class-time or prior to exercise. School may need to make special arrangements for pupils with diabetes if the school has staggered lunchtimes. If a meal or snack is missed, or after strenuous activity, the child may experience a hypoglycaemic episode (a hypo) during which blood glucose level fall too low. Staff in charge of physical education or other physical activity sessions should be aware of the need for children with diabetes to have glucose tablets or a sugary drink to hand.

Staff should be aware that the following symptoms, either individually or combined, may be indicators of low blood sugar – a **hypoglycaemic reaction** (hypo) in a child with diabetes:

- hunger
- sweating
- drowsiness
- pallor
- glazed eyes
- shaking or trembling
- lack of concentration
- irritability
- headache
- mood changes, especially angry or aggressive behaviour

Insulin pens should be reloaded every day by the family, at home. We will accept insulin vials in school as an absolute emergency supply, where they will be stored in a locked refrigerator.

Each child may experience different symptoms and this should be discussed when drawing up an individual health and risk assessment.

If a child has a hypo, it is very important that the child is not left alone and that a fast acting sugar, such as glucose tablets, a glucose rich gel, or a sugary drink is brought to the child and given immediately. Glucose tablets and gels, if provided, are located within a child's personal diabetic pack.

If after 15 minutes, the levels are back up above 4 then a slower acting starchy food with approx. 10g of carbs e.g, biscuits such a x1 digestive, x1 jaffa cake, x1 custard cream or a glass, should be given to sustain the levels.

If they are still low then repeat the fast acting treatment. If 15 minutes later the levels still haven't risen, call the parent and/or ambulance depending on individual care plam. Always continue with treatment whilst waiting for parent or ambulance.

Hypo and hyper reactions and control measures are advised by a medical professional.

### An ambulance should be called if recovery takes longer than 10-15 minutes or if the person becomes unconscious.

Some children may experience **hyperglycaemia** (high glucose level) and have a greater than usual need to go to the toilet or to drink. Tiredness and weight loss may indicate poor diabetic control, and staff will naturally wish to draw any such signs to the families' attention. If the child is unwell, vomiting or has diarrhoea this can lead to dehydration. If the child is giving off a smell of pear drops or acetone this may be a sign of ketosis and dehydration and the child will need urgent medical attention.

Information and photographs of children with diabetes are in the medical room and other relevant areas and in cohort profile folders as part of care plans.

#### **Training**

Once the school is advised that a child is diabetic then appropriate training and support will be available through the Diabetic Team. Training will be refreshed yearly. Training is provided at 3 different levels. Level 1 and level 2 are suitable for staff who will be monitoring a child with Diabetes. Level 3 training must be undertaken to support the child's administration of insulin. Staff members are not required to undertake Level 3 training if they are not comfortable to do so.

#### **Records**

Records are kept in school and updated by the supporting adults and a second signatory to ensure that the insulin given is correct. Records of insulin given will also be passed to families as part of the child's diabetic pack as well as being recorded on medical devices/ apps.

Medical professionals also request that the school receive a daily diary from the families to advise the school of how many carbohydrates in snacks and lunch boxes, as it is not the responsibility of the school to calculate them individually. If the child has a school lunch, members of school staff will be trained in calculating the number of carbohydrates in the lunches and the size of the portion needed to match the carbohydrates needed at that time.